


## Product Datasheet

### Goat anti-Rabbit IgG (H&L), DyLight® 488 conjugated GRP12256

<b>Species/Host</b>	Goat
<b>Reactivity</b>	Rabbit
<b>Predicted Reactivity</b>	Rabbit IgG Heavy and Light chains (H&L)
<b>Tested Applications</b>	ICC, IF, IHC
<b>Immunogen</b>	Purified Rabbit IgG, whole molecule
<b>Form/Appearance</b>	Lyophilized
<b>Storage</b>	Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified goat IgG
<b>Dilution Range</b>	1 : 50- 1 : 5 000 (ICC), 1 : 20- 1 : 2000 (IHC), 1 : 3000 (IF)



<b>Application Notes</b>	Additional Information: Based in immunoelectrophoresis, this antibody reacts with heavy chains on rabbit IgG and light chains on all rabbit immunoglobulins. No reactivity is observed to non-immunoglobulin rabbit serum proteins based in immunoelectrophoresis. Purity of this antibody is > 95% based on SDS-PAGE. Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. DyLight® 488 has a maximum absorbance at 493 nm; Emax = 518 nm. Background: Goat anti-rabbit IgG, DyLight® 488 Conjugate is a secondary antibody conjugated to DyLight® 488, which binds to all rabbit IgGs in immunological assays. DyLight® is a trademark of Thermo Fisher Scientific, Inc. and its subsidiaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.
--------------------------	---