

## Product Datasheet

### Goat anti-Rabbit IgG (H&L), HRP conjugated, min. cross-reactivity to bovine/Human/goat/mouse/rat IgG GRP12616

<b>Species/Host</b>	Goat
<b>Reactivity</b>	Rabbit
<b>Predicted Reactivity</b>	Rabbit IgG Heavy and Light chains (H&L)
<b>Tested Applications</b>	ELISA, WB, IHC
<b>Immunogen</b>	Purified Rabbit IgG, whole molecule (H&L chains)
<b>Form/Appearance</b>	Lyophilized
<b>Storage</b>	Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified goat IgG
<b>Dilution Range</b>	1 : 200- 1 : 5 000 (ELISA), 1 : 500- 1 : 5 000 (IHC), 1 : 200- 1 : 5 000 (WB)
<b>Application Notes</b>	Additional Information: No reactivity is observed to non-immunoglobulin rabbit serum proteins based in immunoelectrophoresis. No reactivity is observed to bovine/goat/human/mouse or rat IgG based on immunoelectrophoresis. No reactivity to non-immunoglobulin rabbit serum proteins. Antibody is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.1 % (v/v) Kathon CG is added as preservative. Use of sodium azide will inhibit enzyme activity of horseradish peroxidase. Background: Goat anti-rabbit IgG is a secondary antibody conjugated to HRP which binds to all rabbit IgGs in immunological assays. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.