

Product Datasheet

Goat anti-Human IgA heavy (alpha chain), ALP conjugated, min. cross-reactivity to bovine/mouse/rabbit serum GRP12626

Species/Host	Goat
Reactivity	Human
Predicted Reactivity	Human IgA, Heavy alpha chains
Tested Applications	ELISA, ICC, WB, IHC
Immunogen	purified human IgA, alpha chain
Form/Appearance	Liquid
Storage	Non-diluted antibody is stable for 4 years at 2-8°C. For storage at -20°C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified goat IgG
Dilution Range	1 : 500-1 : 2000 (ELISA) , 1 : 50-1 : 5000 (ICC), 1 : 20 -1 : 2000 (IHC), 1 : 500-1 : 2000 (WB)
Application Notes	Additional Information: No reactivity is observed to light chains or non-IgA human serum proteins based in immunoelectrophoresis no reactivity to bovine, mouse or rabbit serum proteins based on immunoelectrophoresis AP conjugate is supplied in 30 mM Triethanolamine, pH 7.2, 5 mM Magnesium Chloride, 0.1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) of sodium azide is added as preservative Background: Goat anti-human IgA heavy (alpha chain) is a secondary antibody conjugated to AP (Alkaline phosphatase) which binds to human IgA heavy, alpha chains in immunological assays.