

Product Datasheet

Goat anti-Bovine IgG (H&L), FITC conjugated GRP12991

Species/Host	Goat
Reactivity	Bovine
Tested Applications	FC, IL
Immunogen	Purified Bovine IgG, whole molecule
Form/Appearance	Liquid
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Uniprot ID	DAA31645
Dilution Range	1 : 20-1 : 2000 (Flow cys), (IL). The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Bovine IgG (H&L). Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">· heavy (γ) chains on bovine IgG· light chains on all bovine immunoglobulins. <p>Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none">· non-immunoglobulin bovine serum immunoglobulins. <p>Background: Rabbit anti-Bovine IgG (H&L), FITC conjugated, is a secondary antibody conjugated to FITC which binds to Bovine IgG (H&L) in immunological assays. Amax = 494 nm; Emax = 518 nm.</p>