

Product Datasheet

Goat anti-Bovine IgG (H&L), TRITC conjugated GRP12994

Species/Host	Goat
Reactivity	Bovine
Tested Applications	FC, IL
Immunogen	Purified Bovine IgG, whole molecule
Form/Appearance	Lyophilized
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Uniprot ID	DAA31645
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	<p>Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Bovine IgG (H&L). Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on bovine IgG• light chains on all bovine immunoglobulins <p>Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none">• non-immunoglobulin bovine serum immunoglobulins <p>Background: Goat anti-Bovine IgG (H&L), TRITC conjugated, is a secondary antibody conjugated to TRITC which binds to Bovine IgG (H&L) in immunological assays. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.</p>