

Product Datasheet

Goat anti-Cat IgG (H&L), TRITC conjugated GRP12997

| | |
|----------------------------|--|
| Species/Host | Goat |
| Reactivity | Cat |
| Tested Applications | FC, IL |
| Immunogen | Purified Cat IgG, whole molecule |
| Form/Appearance | Lyophilized |
| Storage | Store at 2-8 °C. |
| Note | For research use only. |
| Clonality | Polyclonal |
| Purity | Affinity purified |
| Dilution Range | 1 : 20-1 : 2000 (Flow cyt), (IL) |
| Application Notes | <p>Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Cat IgG (H&L). Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on cat IgG• light chains on all cat immunoglobulins <p>Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none">• non-immunoglobulin cat serum immunoglobulins <p>Background: Goat anti-Cat IgG (H&L) is a secondary antibody, conjugated to TRITC, which binds to Cat IgG (H&L) in immunological assays. Reconstitution: For reconstitution add 0.55 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.</p> |