

## Product Datasheet

### Goat anti-Human IgG Fc, min. cross-reactivity to human IgA + IgM, bovine, horse, mouse or rabbit serum proteins GRP13039

<b>Species/Host</b>	Goat
<b>Reactivity</b>	Human
<b>Immunogen</b>	Purified Human IgG, Fc fragment
<b>Form/Appearance</b>	Liquid
<b>Storage</b>	Store at 2-8 °C.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified
<b>Uniprot ID</b>	O75015
<b>Dilution Range</b>	The optimal working dilution should be determined by the investigator.

**Application Notes** Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Human IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy (γ) chains on human IgG. Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgA or IgM • serum proteins from bovine, horse, mouse, or rabbit. Background: Goat anti-Human IgG Fc, min. cross-reactivity to bovine, horse, mouse or rabbit serum proteins or mouse IgG1 is a secondary antibody which binds to human IgG Fc in immunological assays.