

## Product Datasheet

# Donkey anti-Mouse IgG (H&L), F(ab)'2 fragment, Biotin conjugated, min. cross-reactivity to bovine, chicken, goat, guinea pig, hamster, horse, human, rabbit, rat or sheep IgG GRP13075

<b>Species/Host</b>	Donkey
<b>Reactivity</b>	Mouse
<b>Tested Applications</b>	FC, IL
<b>Immunogen</b>	Purified Mouse IgG, whole molecule
<b>Form/Appearance</b>	Liquid
<b>Storage</b>	Store at 2-8 °C.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified
<b>Dilution Range</b>	1 : 20-1 : 2000 (Flow cyt), (IL)
<b>Application Notes</b>	<p>Additional Information: Antibody purity is &gt; 90% based on SDS-PAGE Small amounts of intact IgG may be present. Affinity purified using solid phase Mouse IgG .Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none"><li>• heavy (γ) chains on mouse IgG</li><li>• light chains on all mouse immunoglobulins</li></ul> <p>.Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none"><li>• non-immunoglobulin mouse serum immunoglobulins</li><li>• IgG from bovine, chicken, goat, guinea pig, hamster, horse, human, rabbit, rat or sheep .</li></ul> <p>Background: Donkey anti-Mouse IgG (H&amp;L), F(ab)'2 fragment, Biotin conjugate, min. cross-reactivity to bovine, chicken, goat, guinea pig, hamster, horse, human, rabbit, rat or sheep IgG is a secondary antibody conjugated to biotin, which binds to mouse IgG (H&amp;L), F(ab)'2 Fragment in immunological assays.</p>