

## Product Datasheet

### Goat anti-Mouse IgG Fc, F(ab)<sup>'</sup>2 Fragment GRP13110

<b>Species/Host</b>	Goat
<b>Reactivity</b>	Mouse
<b>Tested Applications</b>	ELISA, IP, WB, IHC, IDA
<b>Immunogen</b>	Purified Mouse IgG Fc
<b>Form/Appearance</b>	Liquid
<b>Storage</b>	Store at 2 - 8 °C.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified
<b>Dilution Range</b>	The optimal working dilution should be determined by the investigator.
<b>Application Notes</b>	Additional Information: Antibody purity is > 90 % based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Mouse IgG. F(ab) <sup>'</sup> 2 fragment was prepared by pepsin digestion. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05 % (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: heavy (γ) chains on Mouse IgG. Based on IEP, no reactivity is observed to: light chains on all Mouse immunoglobulins, non-immunoglobulin Mouse serum proteins, Mouse IgG, F(ab) <sup>'</sup> 2 fragment, serum proteins from Bovine, Horse, or Human. Background: Goat anti-Mouse IgG Fc, F(ab) <sup>'</sup> 2 Fragment is a secondary antibody which binds to mouse IgG Fc F(ab) <sup>'</sup> 2 fragment in immunological assays.