

Product Datasheet

Goat anti-Mouse IgG Fc, F(ab)[']2 Fragment GRP13110

Species/Host	Goat
Reactivity	Mouse
Tested Applications	ELISA, IP, WB, IHC, IDA
Immunogen	Purified Mouse IgG Fc
Form/Appearance	Liquid
Storage	Store at 2 - 8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	Additional Information: Antibody purity is > 90 % based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Mouse IgG. F(ab) ['] 2 fragment was prepared by pepsin digestion. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05 % (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: heavy (γ) chains on Mouse IgG. Based on IEP, no reactivity is observed to: light chains on all Mouse immunoglobulins, non-immunoglobulin Mouse serum proteins, Mouse IgG, F(ab) ['] 2 fragment, serum proteins from Bovine, Horse, or Human. Background: Goat anti-Mouse IgG Fc, F(ab) ['] 2 Fragment is a secondary antibody which binds to mouse IgG Fc F(ab) ['] 2 fragment in immunological assays.