

## Product Datasheet

### Donkey anti-Rabbit IgG (H&L), F(ab)'2 Fragment GRP13129

<b>Species/Host</b>	Donkey
<b>Reactivity</b>	Rabbit
<b>Immunogen</b>	Purified Rabbit IgG, whole molecule
<b>Form/Appearance</b>	Liquid
<b>Storage</b>	Store at 2-8 °C.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified
<b>Dilution Range</b>	The optimal working dilution should be determined by the investigator.

**Application Notes** Additional Information: Antibody purity is > 90% based on SDS-PAGE Small amounts of intact IgG may be present. Affinity purified using solid phase Rabbit IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy (γ) chains on rabbit IgG • light chains on all rabbit immunoglobulins. Based on IEP, no reactivity is observed to: • non-immunoglobulin rabbit serum proteins. Background: Donkey anti-Rabbit IgG (H&L), F(ab)'2 Fragment is a secondary antibody which binds to rabbit IgG, F(ab)'2 Fragment in immunological assays.