



Product Datasheet

TET1 antibody [N1], N-term GRP64

Description	Dioxygenase that specifically binds methylcytosine (5mC), a minor base in mammalian DNA found in repetitive DNA elements that is crucial for retrotransposon silencing and mammalian development. Catalyzes the conversion of methylcytosine (5mC) to 5-hydroxymethylcytosine (hmC). The clear function of 5-hydroxymethylcytosine (hmC) is still unclear but it may influence chromatin structure and recruit specific factors or may constitute an intermediate component in cytosine demethylation. 5-hydroxymethylcytosine (hmC) is present in ES cells and is enriched in the brain, especially in Purkinje neurons. May play a role in the fetal development of heart, lung and brain.
Species/Host	Rabbit
Reactivity	Human, Mouse
Conjugation	Unconjugated
Tested Applications	ChIP, ICC, IF, IHC-P, WB
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of mouse TET1. The exact sequence is proprietary.
Form/Appearance	Liquid: 1XPBS, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative.
Concentration	1 mg/ml
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Note	For research use only.
Isotype	IgG
Clonality	Polyclonal
Purity	Purified by antigen-affinity chromatography.
Uniprot ID	Q3URK3
Entrez	52463
Dilution Range	WB: 1:500-1:20000, ICC: 1:100-1:1000



TET1 antibody detects TET1 protein by western blot analysis. Non-transfected (-) and mouse TET1-transfected (+, including DDDDK-tag) 293T whole cell extracts (30 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with TET1 antibody (GRP516)



Western blot analysis of 293T cell are mock transfected (lane 1) or with tagged-mTet1 (lane 2) for 24hrs, using either GRP516 antibody. The HRP-conjugated anti-rabbit IgG antibody was used to detect